

2024 S.T. Yau High School Science Award (Asia)

Research Report

The Team

Registration Number: Bio-132

Name of team member: Soha Aggarwal

School: The Cathedral and John Connon School, Mumbai

Country: India

Name of supervising teacher: Aditi Singh

Job Title: Ph.D. Scholar

School: Gautam Buddha University, Greater Noida

Country: India

Title of Research Report Structural and Biophysical studies of β -galactosidase from *Oryza sativa* (Osbg)

Date

14 August 2024

Structural and Biophysical studies of β -galactosidase from *Oryza sativa***(Osbg)****Soha Aggarwal****Abstract**

Beta-galactosidase has a very special space in the history of molecular biology. It proved to be a strong basis for the development of the Jacob and Monod model for gene expression. Here in this work, the enzyme has been isolated from *Oryza sativa*. A simple protocol for extraction and purification of the enzyme was chosen. Anion-exchange chromatography was performed to purify the enzyme. Circular Dichroism gave important information about the α -helix (15.9%) and β -strands (30.4%) in the enzyme. In-silico methods have been implemented to attain the three-dimensional structure of β -galactosidase from *Oryza sativa*. To study the catalytic sites molecular docking has been performed. This work supports the structure-based drug design for developing an analogue in place of non-functional lactase enzymes.

Keywords: beta-galactosidase, gene, molecular biology, molecular docking

Acknowledgement

I would like to express my sincere gratitude to my mentor, Aditi Singh, for her invaluable guidance and support during this research. I would also like to thank Gautam Buddha University, for extending their resources and facilities to me so that I could pursue this research. Finally, I would like to thank my parents for their constant support and encouragement during the entirety of this process, and being my cheerleaders in any venture I pursue.

Research Report

2024 S.T. Yau High School Science Award (Asia)

Commitments on Academic Honesty and Integrity

We hereby declare that we

1. are fully committed to the principle of honesty, integrity and fair play throughout the competition.
2. actually perform the research work ourselves and thus truly understand the content of the work.
3. observe the common standard of academic integrity adopted by most journals and degree theses.
4. have declared all the assistance and contribution we have received from any personnel, agency, institution, etc. for the research work.
5. undertake to avoid getting in touch with assessment panel members in a way that may lead to direct or indirect conflict of interest.
6. undertake to avoid any interaction with assessment panel members that would undermine the neutrality of the panel member and fairness of the assessment process.
7. observe the safety regulations of the laboratory(ies) where the we conduct the experiment(s), if applicable.
8. observe all rules and regulations of the competition.
9. agree that the decision of YHSA(Asia) is final in all matters related to the competition.

We understand and agree that failure to honour the above commitments may lead to disqualification from the competition and/or removal of reward, if applicable; that any unethical deeds, if found, will be disclosed to the school principal of team member(s) and relevant parties if deemed necessary; and that the decision of YHSA(Asia) is final and no appeal will be accepted.

(Signatures of full team below)

Soha Aggarwal

Name of team member: Soha Aggarwal

Aditi Singh

Name of supervising teacher: Aditi Singh

Noted and endorsed by:

(signature)

Name of school principal:

Dr. Sonal Parmar




Table of Contents

Abstract	i
Acknowledgement	ii
Commitments on Academic Honesty and Integrity	iii
1. Introduction	
2. Materials and Methods	
3. Results and discussions	
4. Conclusion	
5. References	

1.0 Introduction

Beta-galactosidase is a hydrolase enzyme that hydrolyses lactose into monosaccharides as glucose and galactose and therefore essential for the digestion of bovine milk [1,2]. The enzyme is secreted by the lacz-gene of the lac operon and deficiency of this gene leads to malabsorption of lactose and ultimately leads to its fermentation in the gut flora [3]. Lactose-intolerant individuals suffer from abdominal pain, diarrhoea, bloating, poor digestion and gastrointestinal symptoms after lactose intake [4]. Although dairy products are a vital source of calcium and vitamin D, people with lactose intolerance have to avoid these dairy products and subsequently causing adverse health outcomes, especially reduced bone mineral density [5,6].

Lactose is a disaccharide formed by the combination of glucose and galactose and its hydrolysis is possible in the presence of lactase, commonly used as β -galactosidase. Generating lactose-free milk can be carried out by introducing the enzyme directly into the milk [7]. In the past various research have been performed in this direction using microbial, animal, plant, fungi and yeast [8,9]. Although bacteria produce lactase enzyme very efficiently, they are not considered a safer source for prior use in food and dairy manufacturing industries due to the risk of microbial contamination. Fungal and yeast sources produce mesophilic lactase enzyme [10]. Different pH optima are required depending on the strains for example acidic and neutral pH for hydrolysis of lactose in whey and milk respectively [11]. Plant-based lactases prove to be at low risk for contamination and also show high enzymic activity and are probably concerned with the metabolism of galactolipids, such as β -D-galactosyl diacylglycerol, universally present in plants and algae [12]. To understand the concerned functions of the enzyme, the catalytic domain of that particular enzyme needs to be explored. The catalytic site is responsible for the activity such as hydrolysis, trans-acetylation, transferase activity and etc [13,14].

In this work, plant-based lactase enzyme has been extracted using a simple protocol and purified through anion-exchange chromatography. Purified enzyme has been obtained and further the activity of the enzyme was checked with ONPG, an analog of lactose. The absorbance of the released o-nitrophenol was recorded at 540 nm. This work reports the three-dimensional structure and catalytic site of the enzyme using molecular docking. There may be a possibility of producing non-functional lactase enzyme by the gene but catalytic sites could help to relate with some substitute that proves to be a novel target in designing any therapeutic treatment for lactose-intolerant people.

2.0 Materials and Methods

2.1 Extraction

Oryza sativa dried seeds were purchased from Etsy, online. Seeds were grounded in a fine powder using a blender. 100 gm of the powder was taken and added in 50 mL of 50mM Tris pH 8. The powder was mixed gently and kept for 20 minutes. Further, the solution was centrifuged at 10,000 rpm for about 20 minutes to collect the supernatant. The supernatant was further proceeded to perform ammonium-sulphate precipitation, initially 30% and then 80%. The overnight incubated solution with 80% salt (ammonium-sulphate) precipitation was centrifuged at 6000 rpm for 10 minutes and pellet was restored at -20°C.

2.2 Purification

For purification, POROS™ XQ (ThermoFisher Scientific) resin were used to perform anion-exchange chromatography [15]. The stored pellet was resuspended in buffer of enzyme (50mM Tris pH 8) and spun down to obtain a clear supernatant. Meanwhile, the column was packed using the resin (POROS™ XQ) and equilibrated with the buffer of protein. The supernatant collected loaded on to the column and allowed to bind with the resin, followed by the washing with the buffer to elute out all the non-specific proteins. As the absorbance at 280nm reaches 0, NaCl gradient was given starting from 0.1M to 0.5M.

2.3 Enzyme Activity

The enzyme assay was carried out using the substrate ONPG (o-nitrophenyl-β-D-galactopyranoside) [16]. The reaction was incubated for 30 minutes in a 1.5 mL Eppendorf with a mixture containing 390μL of phosphate buffer (pH 7.2), 100μL of 5mM ONPG and 10μL of enzyme. After incubation of 30 minutes 500μL of 1M sodium carbonate was added to the mixture to stop the reaction and absorbance was recorded at 420nm using PerkinElmer UV-vis spectrophotometer.

2.4 Circular Dichroism spectroscopic (CDS) studies

CDS measurements of purified β-galactosidase were observed on a JASCO CD Spectrometer. 0.5mg/ml concentration of the enzyme in 50mM Tris pH 8 buffer was used to monitor the structural changes. Absorbance was recorded in far-UV region (190-240nm) using a quartz cuvette with a path length of 1mm.

2.5 3-dimensional structure prediction

The sequence of the enzyme (**Q10NX8 · BGAL6_ORYSJ**) in a FASTA format was obtained from UniProtKB database and used for template-based structure prediction [17]. The 3-Dimensional structure of beta-galactosidase from *Oryza sativa* has been predicted using SWISS-MODEL, tool offered by Expasy: Swiss Institute of Bioinformatics (SIB). Further, PyMOL was used to visualize and analyse the structure. The enzyme has 55.7% similarity with beta-galactosidase from *Solanum lycopersicum* (tomato).

2.6 Molecular Docking

To study the binding sites and the amino acid interaction, molecular docking of beta-galactosidase with NAG (2-acetamido-2-deoxy-beta-D-glucopyranose) was carried out using AutoDockVina. To process the docking, the grid within a spacing of 40 Å × 40 Å × 40 Å centered on the selected flexible residues present in the active site of the enzyme had been chosen. Optimum conformational space had been provided by the grid box to observe rotation to achieve best binding conformation. The complex was visualized on PyMOL [18].

3.0 Results and discussions

3.1 Purification




Figure 1(a) chromatogram for the anion-exchange chromatography. 1(b) purified sample on SDS-PAGE with molecular weight as 80 kDa.

Extracting the enzyme followed by the purification using anion exchange chromatography. The chromatogram was plotted for the fractions collected during the chromatography and the 3rd collected

with 0.3M NaCl gradient had the highest absorbance as shown in figure 1(a). The purity of the sample was depicted on Coomassie blue stained 15% SDS-PAGE and molecular weight was estimated as 80 KDa (figure 1(b)). After concentrating the sample using Amicon filter (30,000 MW cut-off) the concentration was found to be 5mg/ml.

3.1 Activity and secondary structural studies (CD)




Figure 2(a) enzyme-substrate catalysis depicting the maximum activity of the enzyme. (b) CD spectra of the enzyme in far-UV region.

The hydrolysing activity of the enzyme was measured using colorimetric assay. The enzyme was found catalytically active which hydrolyzes ONPG into o-nitrophenol. The rate of catalytic reaction increased linearly with time but did not reach V_{max} in 30 minutes (figure 2(a)). CD spectroscopy far-UV measurements of the sample depicted 15.9% α -helices and 30.4% β -strands as analyzed through K2D3 online server for data analysis (figure 2(b)) [19].

3.2 Structure of Osbg




Figure 3 three-dimensional structure of Osbg .

The bioinformatics studies suggested that the Osbg depicts highest similarity with the crystal structure of tbg (PDB ID: 6ik5) with sequence identity of 55.5%. The secondary structure contains 40 strands, 20 helices and 59 coiled regions (figure 3) as estimated by PDBsum and verifiable with the data obtained from CD analysis. The LG score of the model was found to be 4.877 which brings it in the list of good quality models. The results suggest that the 3-D model for Osbg is reliable. On the basis Ramachandran plot, 87.9% of the residues were found in the most favored region, 11.6 % in the additionally allowed region and 0.3%residues in the disallowed region.

3.3 NAG binding site




Figure 4 active participation of the residues with the ligand(NAG) through molecular docking.

NAG was docked on Osbg in order to obtain the recognition and position of catalytic site of the enzyme. As shown in figure 4, the ligand was located and stabilized by attractive interactions with LEU 516, ASN 158, MET 159, GLU 163, VAL 96, and PHE 152 residues in the substrate binding site of the enzyme.

4. Conclusion

Osbg enzyme was extracted from oryza sativa and purified to homogeneity using chromatographic technique. The CD measurements depict about the stable three-dimensional structure, predominantly with the presence β -strands. Osbg forms a very stable complex with NAG and describes the active participation of the catalytic residues of the enzyme. The in-silico studies suggested that the structure is similar to the B-gal from *S. lycopersium*. The whole study proves to be a unique possibility for the people where the enzyme is produced non-functionally. The active site of the enzyme provides the information about the target for futuristic drug developments.

References:

- [1] Shukla, T. P., & Wierzbicki, L. E. (1975). Beta-galactosidase technology: A solution to the lactose problem. *Critical Reviews in Food Science & Nutrition*, 5(3), 325-356.
- [2] Richmond, M. L., Gray, J. I., & Stine, C. M. (1981). Beta-galactosidase: review of recent research related to technological application, nutritional concerns, and immobilization. *Journal of dairy science*, 64(9), 1759-1771.
- [3] Mason, J. M., Hackett, R. H., & Setlow, P. (1988). Regulation of expression of genes coding for small, acid-soluble proteins of *Bacillus subtilis* spores: studies using lacZ gene fusions. *Journal of bacteriology*, 170(1), 239-244.
- [4] Swagerty Jr, D. L., Walling, A. D., & Klein, R. M. (2002). Lactose intolerance. *American family physician*, 65(9), 1845-1851.
- [5] Kudlacek, S., Freudenthaler, O., Weissböeck, H., Schneider, B., & Willvonseder, R. (2002). Lactose intolerance: a risk factor for reduced bone mineral density and vertebral fractures?. *Journal of gastroenterology*, 37, 1014-1019.
- [6] Hamilton, N. K., Ojo, O., & Adegboye, A. R. A. (2020). The effect of self-reported lactose intolerance and dairy consumption on bone mineral density among American hip arthroplasty patients: A cross-sectional study. *International Journal of Environmental Research and Public Health*, 17(19), 7182.
- [7] Xavier, J. R., Ramana, K. V., & Sharma, R. K. (2018). β -galactosidase: Biotechnological applications in food processing. *Journal of Food Biochemistry*, 42(5), e12564.
- [8] Kocabaş, D. S., Lyne, J., & Ustunol, Z. (2022). Hydrolytic enzymes in the dairy industry: Applications, market and future perspectives. *Trends in Food Science & Technology*, 119, 467-475.
- [9] Kalathinathan, P., Sain, A., Pulicherla, K., & Kodiveri Muthukaliannan, G. (2023). A Review on the Various Sources of β -Galactosidase and Its Lactose Hydrolysis Property. *Current Microbiology*, 80(4), 122.
- [10] Soydan, M. (2006). Production of thermostable beta-galactosidase from tyhermophilic fungi for use in low-lactose milk production (Master's thesis, Middle East Technical University).
- [11] Shi, X., Wu, D., Xu, Y., & Yu, X. (2022). Engineering the optimum pH of β -galactosidase from *Aspergillus oryzae* for efficient hydrolysis of lactose. *Journal of Dairy Science*, 105(6), 4772-4782.
- [12] Qi, Y., Yamauchi, Y., Ling, J., Kawano, N., Li, D., & Tanaka, K. (2004). Cloning of a putative monogalactosyldiacylglycerol synthase gene from rice (*Oryza sativa* L.) plants and its expression in response to submergence and other stresses. *Planta*, 219, 450-458.
- [13] Andres, H. H., Kolb, H. J., Schreiber, R. J., & Weiss, L. (1983). Characterization of the active site, substrate specificity and kinetic properties of acetyl-CoA: arylamine N-acetyltransferase from pigeon liver. *Biochimica et Biophysica Acta (BBA)-Protein Structure and Molecular Enzymology*, 746(3), 193-201.
- [14] Zou, S., Li, X., Huang, Y., Zhang, B., Tang, H., Xue, Y., & Zheng, Y. (2023). Properties and biotechnological applications of microbial deacetylase. *Applied Microbiology and Biotechnology*, 1-20.
- [15] Lee, Y. C. (1990). High-performance anion-exchange chromatography for carbohydrate analysis. *Analytical biochemistry*, 189(2), 151-162.
- [16] Ladero, M., Santos, A., García, J. L., & García-Ochoa, F. (2001). Activity over lactose and ONPG of a genetically engineered β -galactosidase from *Escherichia coli* in solution and immobilized: Kinetic modelling. *Enzyme and Microbial technology*, 29(2-3), 181-193.
- [17] Martin, A. C. (2005). Mapping PDB chains to UniProtKB entries. *Bioinformatics*, 21(23), 4297-4301.

[18] Yuan, S., Chan, H. S., & Hu, Z. (2017). Using PyMOL as a platform for computational drug design. *Wiley Interdisciplinary Reviews: Computational Molecular Science*, 7(2), e1298.

[19] Louis-Jeune, C., Andrade-Navarro, M. A., & Perez-Iratxeta, C. (2012). Prediction of protein secondary structure from circular dichroism using theoretically derived spectra. *Proteins: Structure, Function, and Bioinformatics*, 80(2), 374-381.